

# Sacroiliac Joint Injection

## Patient Information Sheet

### 1. What is a sacroiliac joint injection?

A sacroiliac joint injection is an injection of local anesthetic and steroid deposited into the sacroiliac joint. At the lower end of the spine, just below the lumbar spine lies the sacrum. The sacrum is a triangular-shaped bone that is actually formed by the fusion of several vertebrae during development. The sacroiliac (SI) joint sits between the sacrum and the iliac bone (also called the ilium.) This is why it is called the sacroiliac joint. The goal of a sacroiliac joint injection is to provide pain relief by reducing the inflammation of the joint. A sacroiliac joint injection will not correct the preexisting medical problem (i.e. bone spurs, arthritis, etc.) but may improve the level of pain. It is not unusual for someone to need more than one injection to get long term benefit. The injections are done in a series of three injections about 2-4 weeks apart if needed. If the pain significantly improves, no further injection is needed unless the pain begins to come back.

**Note: The procedure can not be performed if you have an active infection, flu, cold, fever, very high blood pressure, or if you are on blood thinners. Please make your doctor aware of any of these conditions. This is for your safety!**

### 2. What are the risks of the procedure?

Risks are remote but include bleeding, infection, nerve injury, and allergic reaction to the medication.

Some short-term side effects may occur. If local anesthetic spreads to nearby nerves you may have weakness or numbness that can last for 1 – 2 hours. If this happens you will have to stay in the Murphy Pain Center until this resolves. You may have increased pain for a few days after the injection. Diabetics may have short-term elevation of blood sugars. People prone to fluid retention may have increased fluid retention for 1 – 2 weeks.

### 3. Will the injection hurt a lot?

Most people say the stinging/burning of the numbing medicine is the most uncomfortable part of the procedure though every person's response to any procedure is different.

### 4. What happens during the actual procedure?

After signing a consent form and checking your blood pressure the procedure will be done in the fluoroscopy (x-ray) room with you lying on your stomach. The lower back is then cleansed with an antiseptic soap and a drape is placed. The needle is then advanced into the sacroiliac joint. Pressure is the usual sensation felt. Once the needle is in the sacroiliac joint, the medication is infused, and the procedure is complete. Your skin will be cleansed and a bandage will be applied. (The bandage can be removed when you get home). Your vital signs will be checked. You will be escorted to the recovery area where you will be offered something to eat and drink. You will be monitored in the recovery area for 15 minutes. Your pain level will then be re-assessed and you will be discharged to leave with your ride.

### 5. How will I feel after the injection?

Most people don't feel any different immediately after the injection though the pain may be temporarily improved or worsened. The steroid takes two or three days to start to have an effect in most people and peaks in about two weeks. Therefore, it may be a while before you feel a change in you pain.

Some local tenderness may be experienced for a couple of days after the injection. You will be given instructions on the use of ice and heat at your discharge to help minimize this discomfort. You may take your usual pain medications as well after the injection.

**6. Will I have any restrictions on the day of the procedure?**

**Do not eat or drink** for 6 hours prior to the procedure. You may take your medications for blood pressure or heart with a sip of water. You may also take your pain medication with a sip of water if you are able to tolerate it on an empty stomach. **If you are DIABETIC or take BLOOD THINNERS, you will need to contact the office prior to your procedure** for instructions, unless you received instructions when you scheduled the appointment. Be aware that some types of blood thinners must be stopped several days prior to your injection which will require the approval of your prescribing physician.

You may not drive for the remainder of the day after your procedure. An adult must be present to drive you home or to go with you in a taxi or on public transportation. **The procedure will be canceled if you don't have a responsible adult with you!!! This is for your safety.**

No heat is to be used in the injected area for the remainder of the day.

No tub bath or soaking in water (i.e. pool, jacuzzi, etc.) for the remainder of the day.

You may return to eating and drinking as usual after your procedure.

**7. For what reasons should I call the Murphy Pain Center after the injection?**

If you experience severe back pain, new numbness or weakness of your legs, loss of control of your bowels or bladder, or signs of infection in the area of the injection, you should call the Murphy Pain Center right away at 502-736-3636

References:

Harvard Medical School-Brigham and Women's Hospital Pain Management Center.

<http://www.hmcnet.harvard.edu/brighampain/faqs/epidural.html>

A Patient's Guide to Sacroiliac Joint Injections

<http://www.eorthopod.com/content/sacroiliac-joint-injections>